

Contents lists available at ScienceDirect

Systems & Control Letters



journal homepage: www.elsevier.com/locate/sysconle

Global linearization of asymptotically stable systems without hyperbolicity

Matthew D. Kvalheim^a, Eduardo D. Sontag^b,*

^a Department of Mathematics and Statistics, University of Maryland, Baltimore County, MD, USA
^b Departments of Electrical and Computer Engineering and Bioengineering, and affiliate of Departments of Mathematics and Chemical Engineering, Northeastern University, Boston, MA, USA

ARTICLE INFO

Keywords: Linearization Hartman-Grobman theorem

ABSTRACT

We give a proof of an extension of the Hartman-Grobman theorem to nonhyperbolic but asymptotically stable equilibria of vector fields. Moreover, the linearizing topological conjugacy is (i) defined on the entire basin of attraction if the vector field is complete, and (ii) a $C^{k\geq 1}$ -diffeomorphism on the complement of the equilibrium if the vector field is C^k and the underlying space is not 5-dimensional. We also show that the C^k statement in the 5-dimensional case is equivalent to the 4-dimensional smooth Poincaré conjecture.

1. Introduction

Consider a nonlinear system of ordinary differential equations

 $\dot{x}(t) = f(x(t)),\tag{1}$

where *f* is a vector field on an *n*-dimensional manifold *M*. If $f \in C^1$ and $x_* \in M$ is a hyperbolic equilibrium for *f*, the Hartman-Grobman theorem guarantees existence of continuous local coordinates on a neighborhood of x_* in which the *nonlinear* dynamics (1) become *linear* [1,2].

In this paper, we provide a proof of an extension of the Hartman-Grobman theorem to nonhyperbolic but asymptotically stable equilibria. We also assume only that f is continuous and uniquely integrable (e.g. locally Lipschitz). But if additionally $f \in C^k$ and $n \neq 5$, we construct continuous linearizing coordinates that are C^k away from x_* . Finally, if f is complete, we prove the existence of globally linearizing coordinates on the entire basin of attraction of x_* . In fact, the linear dynamics can be taken to be $\dot{y} = -y$, in which case the coordinates transform the nonlinear dynamics into the negative gradient flow of the convex function $y \mapsto ||y||^2/2$.

To our knowledge, the first trace of these extensions appeared in work of Coleman, who proved a theorem equivalent to our local result for k = 0 assuming x_* has a Lyapunov function with level sets homeomorphic to spheres [3,4] (see [5, p. 247]). Our proof of the global result fills in details of one previously sketched by Grüne, the second author, and Wirth for k = 0 and $n \neq 4$, and for general kand $n \neq 4,5$ [6, remark in p. 133]. Using the same techniques and Perelman's solution to the 3-dimensional Poincaré conjecture [7–9], we extend this global result to all n for k = 0, and to $n \neq 5$ for general k. In fact, we prove that whether the $C^{k\geq 1}$ linearization results hold for n = 5 is *equivalent* to the 4-dimensional smooth Poincaré conjecture (still open). Jongeneel recently proved other interesting stability results using Perelman's solution [10]. We also note connections to classical dynamical systems results showing that systems with globally asymptotically stable equilibria admit C^0 Lyapunov functions V with exponential decrease $V(x(t)) = e^{-t}V(x(0))$ [11, Chapter V.2]. Furthermore, in the special case of linear systems, it has long been known that each linear system is equivalent, under a continuous coordinate change, to the system $\dot{y} = -y$ of the same dimension [12].

The local linearization result is Theorem 1, and the global result is Theorem 2. While some techniques in the hyperbolic case establish global linearizations by extending local ones [13–15], in this paper we prove the global result directly (and under no hyperbolicity assumptions) and the local result as a consequence. In contrast to the classical hyperbolic results, we assume stability, since otherwise there are no natural global Lyapunov functions to guide the constructions.

It seems worth noting that Theorems 1, 2 give new existence results for targets of algorithms like extended Dynamic Mode Decomposition [16] studied by applied Koopman operator theorists [17–19]. Such algorithms seek to compute $N \ge \dim M$ linearizing "observables" through which nonlinear systems appear linear. Theorems 1, 2 (see also Remark 2) give existence results in the case $N = \dim M$, a case of arguably practical importance [20] complementing recent (non)existence results for the case $N > \dim M$ [21–26].

2. Results

We begin with the local linearization result, whose statement refers to the initial value problem

$$\dot{x}(t) = f(x(t)), \quad x(0) = x_0.$$
 (2)

* Corresponding author. *E-mail addresses:* kvalheim@umbc.edu (M.D. Kvalheim), sontag@sontaglab.org, e.sontag@northestern.edu (E.D. Sontag).

https://doi.org/10.1016/j.sysconle.2025.106163

Received 16 February 2025; Received in revised form 14 May 2025; Accepted 25 May 2025 Available online 13 June 2025

^{0167-6911/© 2025} The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

Theorem 1. Let x_* be an asymptotically stable equilibrium for a uniquely integrable continuous vector field f on an *n*-dimensional C^{∞} manifold M. There is an open neighborhood $U \subset M$ of x_* such that, for any Hurwitz matrix $A \in \mathbb{R}^{n \times n}$, there is a topological embedding $h: U \to \mathbb{R}^n$ such that, for all $x_0 \in U$, the maximal solution of (2) satisfies $x(t) \in U$ and

$$x(t) = h^{-1}(e^{At}h(x_0))$$
 for all $t \ge 0.$ (3)

Moreover, if $n \neq 5$ and $f \in C^k$ with $k \in \mathbb{N}_{\geq 1} \cup \{\infty\}$, then there exists such an *h* that additionally restricts to a C^k embedding $U \setminus \{x_*\} \to \mathbb{R}^n \setminus \{0\}$.

An advantage of Theorem 1 is that it does not assume completeness of f. On the other hand, the global linearization result below is formulated more optimally for smoothness, since the vector field generating a C^k flow need only be C^{k-1} in general.

Theorem 2. Let x_* be an asymptotically stable equilibrium with basin of attraction *B* for the flow Φ of a complete uniquely integrable continuous vector field on an *n*-dimensional C^{∞} manifold *M*. For any Hurwitz matrix $A \in \mathbb{R}^{n \times n}$ there is a homeomorphism $h : B \to \mathbb{R}^n$ satisfying

$$\Phi^t|_B = h^{-1} \circ e^{At} \circ h \quad \text{for all} \quad t \in \mathbb{R}.$$
(4)

Moreover, if $n \neq 5$ and $\Phi \in C^k$ with $k \in \mathbb{N}_{\geq 1} \cup \{\infty\}$, then there exists such an *h* that additionally restricts to a C^k -diffeomorphism $B \setminus \{x_*\} \to \mathbb{R}^n \setminus \{0\}$.

Remark 1. In particular, consider $A = -I_{n \times n}$ in the above theorems. Then as noted in Section 1, *h* transforms the nonlinear dynamics into the negative gradient flow of the dynamics $\dot{y} = -y$ of the convex function $y \mapsto ||y||^2/2$. This is distinct from the fact that a Riemannian metric always exists making the vector field a gradient on the complement of the equilibrium within the basin of attraction [27, Thm 1] in all dimensions (but for $n \neq 5$ this directly follows from the above theorems). The same explanation shows that *h* transforms the nonlinear dynamics into a "contractive system" [28]. There are generally no explicit expressions for the transformation *h*, except in very simple cases. For example, the one-dimensional system $\dot{x} = -x^3$ is transformed into $\dot{y} = -y$ by means of the transformation $y = h(x) = e^{-\frac{1}{2x^2}}$.

Remark 2. By choosing *A* to be diagonal, the conclusion of Theorem 2 furnishes *n*-tuples (ψ_1, \ldots, ψ_n) of continuous real eigenfunctions of the Koopman operator such that the mapping (ψ_1, \ldots, ψ_n) : $B \to \mathbb{R}^n$ is a homeomorphism. Indeed, we may consider the functions $\psi_i = h_i$, where *h* is the homeomorphism from our theorem statement with components h_1, \ldots, h_n .

Remark 3. If the equilibrium x_* is hyperbolic, then the dimensional restriction $n \neq 5$ in Theorems 1, 2 is not needed since then there exists a smooth Lyapunov function having nondegenerate quadratic (Morse) singularity at x_* . The nonzero level sets of such a Lyapunov function are diffeomorphic to the standard sphere of the appropriate dimension, so any such level set can be used instead of *L* in the proof of Theorem 2.

Remark 4. The proofs of Theorems 1 and 2 require only that the vector field *f* is C^k on the complement of $\{x_*\}$ and the flow Φ is C^k on $\mathbb{R} \times B \setminus \{x_*\}$.

The final result establishes the mentioned relationship to the 4dimensional smooth Poincaré conjecture. This conjecture asserts that every 4-dimensional C^{∞} manifold homotopy equivalent to the 4-sphere is diffeomorphic to the 4-sphere.

Proposition 1. Fix $k \in \mathbb{N}_{\geq 1} \cup \{\infty\}$. The 4-dimensional smooth Poincaré conjecture is true if and only if the C^k statement of Theorem 1 (or Theorem 2) is true for n = 5.

3. Proofs

We first assume Theorem 2 to give a short proof of Theorem 1.

Proof of Theorem 1. Fix any Hurwitz matrix $A \in \mathbb{R}^{n \times n}$. Let $\psi : M \to [0, \infty)$ be a C^{∞} function equal to 1 on a neighborhood U_0 of x_* and equal to zero outside of a compact subset of M. Then ψf is a complete uniquely integrable continuous vector field generating a flow Φ , x_* is asymptotically stable for Φ , and $\Phi \in C^k$ if $f \in C^k$. According to Wilson, there is a proper strict C^{∞} Lyapunov function $V : B \to [0, \infty)$ for x_* and Φ [29, Thm 3.2] (see also [30, Sec. 6]). Theorem 2 furnishes a homeomorphism $h_0 : B \to \mathbb{R}^n$ that satisfies (4) and, if $n \neq 5$, restricts to a C^k -diffeomorphism $B \setminus \{x_*\} \to \mathbb{R}^n \setminus \{0\}$. Let c > 0 be sufficiently small that $U := V^{-1}([0, c))$ is contained in $U_0 \cap B$. Then $h := h_0|_U : U \to \mathbb{R}^n$ is the desired embedding.

We now prove Theorem 2 (without assuming Theorem 1).

Proof of Theorem 2. It suffices to find $h: B \to \mathbb{R}^n$ satisfying (4) for A = -I. Indeed, repeating the proof with *B* and *f* replaced by \mathbb{R}^n and $\tilde{f}(x) = Ax$ then yields a corresponding transformation $\tilde{h}: \mathbb{R}^n \to \mathbb{R}^n$, so the composition $\tilde{h}^{-1} \circ h: B \to \mathbb{R}^n$ satisfies (4) for general *A*.

Let $V: B \to [0, \infty)$ be a proper strict C^{∞} Lyapunov function for x_* and Φ ([29, Thm 3.2], [30, Sec. 6]). Fix any c > 0 and note that $V^{-1}([0, c])$ is a contractible compact C^{∞} embedded submanifold with codimension-1 boundary $L := V^{-1}(c)$ homotopy equivalent to the (n-1)-sphere $\mathbb{S}^{n-1} := \{y \in \mathbb{R}^n : ||y|| = 1\}$ [31, pp. 326–327]. Thus, if $n \neq 5$, there is a C^{∞} -diffeomorphism $P: L \to \mathbb{S}^{n-1}$ according to classical facts for n = 2 [32, Appendix] and n = 3 [33, Thm 9.3.11], Perelman for n = 4 [7–9] (see [34, Cor. 0.2]), and Smale for $n \ge 6$ [35, Thm 5.1]. If n = 5 there is still a homeomorphism $P: L \to \mathbb{S}^{n-1}$ according to Freedman [36, Thm 1.6].

Since for each $x \in B \setminus \{x_*\}$ the trajectory $t \mapsto \Phi^t(x)$ converges to x_* and crosses *L* exactly once and transversely, the map $\mathbb{R} \times L \to B \setminus \{x_*\}$ defined by $(t, x) \mapsto \Phi^t(x)$ is a homeomorphism and C^k -diffeomorphism if $\Phi \in C^k$, with inverse $g = (\tau, \rho) : B \setminus \{x_*\} \to \mathbb{R} \times L$ satisfying $\tau(x) \to -\infty$ as $x \to x_*$ (cf. [31, p. 327]). Thus $h : B \to \mathbb{R}^n$ defined by $h(x_*) = 0$ and

$$h(x) = e^{\tau(x)} P(\rho(x))$$

is a homeomorphism that, if $n \neq 5$ and $\Phi \in C^k$, restricts to a C^k diffeomorphism $B \setminus \{x_*\} \rightarrow \mathbb{R}^n \setminus \{0\}$. Finally, *h* satisfies (4) with $A = -I_{n \times n}$ since $\rho \circ \Phi^t |_{B \setminus \{x_*\}} = \rho$ and $\tau \circ \Phi^t |_{B \setminus \{x_*\}} = \tau - t$. \Box

Finally, we prove Proposition 1.

Proof of Proposition 1. Assume that the 4-dimensional smooth Poincaré conjecture is true. Then $P: L \to \mathbb{S}^4$ in the proof of Theorem 2 can be taken to be a diffeomorphism when n = 5. With this fact, repeating the proofs of Theorem 2 and Theorem 1 verbatim show that their C^k statements are true for n = 5.

Conversely, assume that the C^k statement of Theorem 1 (or Theorem 2) is true for n = 5 and $A = -I_{n \times n}$. Let *L* be any 4-dimensional C^{∞} homotopy sphere. According to Hirsch [37, Thm 2], there is a C^{∞} function $V : \mathbb{S}^5 \to [0, 1]$ having only two critical points, a maximum $N = V^{-1}(1)$ and minimum $S = V^{-1}(0)$, such that there is a diffeomorphism $P_c : L \to V^{-1}(c)$ for any $c \in (0, 1)$.

Equip \mathbb{S}^5 with a Riemannian metric and let Φ be the flow of the complete C^{∞} vector field $f = -\nabla V$ on \mathbb{S}^5 . Observe that $S \in \mathbb{S}^5$ is asymptotically stable with basin of attraction $B := \mathbb{S}^5 \setminus \{N\}$ and that f points inward at the boundary $V^{-1}(c)$ of $V^{-1}([0, c])$ for any $c \in (0, 1)$. Thus, Theorem 1 (or Theorem 2) furnishes an open neighborhood $U \subset \mathbb{S}^5$ of S and a C^k embedding $h: U \to \mathbb{R}^5$ such that, for any c > 0 small enough that $V^{-1}(c) \subset U$, h diffeomorphically maps $V^{-1}(c)$ onto a C^k embedded submanifold $L_c \subset \mathbb{R}^5$ intersecting every line through the origin in a single point and transversely. Fixing such a small c > 0

and letting $\rho : \mathbb{R}^5 \setminus \{0\} \to \mathbb{S}^4$ be the straight line retraction $\rho(y) = \frac{y}{\|y\|}$, it follows that

$$L \xrightarrow{P_c} V^{-1}(c) \xrightarrow{h} L_c \xrightarrow{\rho} \mathbb{S}^4$$

is a well-defined C^k -diffeomorphism. Since L is a C^{∞} manifold C^k -diffeomorphic to \mathbb{S}^4 , L is C^{∞} -diffeomorphic to \mathbb{S}^4 [33, Thm 2.2.7].

Additional comments

The paper [6] dealt, more generally, with systems $\dot{x} = f(x, u)$, where *u* denotes an input or disturbance. That paper showed that the input to state stability property [38] is equivalent, under similar global changes of variables, to finiteness of the L^2 norm (" H_{∞} gain") of the input/state operator $(x(0), u(\cdot)) \mapsto x(\cdot)$. As remarked in [6], however, the generalization of linearization constructions to systems with disturbances is not immediate. We thus leave open the study of such extensions.

We also note the relationship between the work reported here and areas of current machine learning research. In [39, Chapter 6] one finds a discussion of "autoencoder" deep neural networks for the numerical approximation of Hartman-Grobman-like conjugacies for linearizations and Koopman eigenvalues. Our results, especially in combination with new theoretical results about the existence of such autoencoders [40], contributes to the theoretical foundation for such studies.

Finally, the proof of Theorem 2 is closely related to the proof of [6, Prop. 1], which can be viewed as a (global) extension of the Morse lemma [41] (see [42, Lem. 2.2]) to local minima of non-Morse functions. Incorporating Perelman's result as in the proof of Theorem 2, the proof of [6, Prop. 1] can otherwise be repeated verbatim to prove Proposition 2 below. It extends the two-part statement [6, Prop. 1] by removing the hypothesis " $n \neq 4$ " from the first part and relaxing the hypothesis " $n \neq 4$, 5" from the second part to " $n \neq 5$ ", which reflects Perelman's result. (It also contains the superficial extension of replacing the domain \mathbb{R}^n of V with a C^{∞} manifold M, but the hypotheses imply that M is diffeomorphic to \mathbb{R}^n .) Proposition 2 is a global statement, but it readily implies a local statement in a manner similar to the implication of Theorem 1 by Theorem 2.

For the following statement, a class \mathcal{K}_{∞} function is a strictly increasing and continuous function $\gamma : [0, \infty) \to [0, \infty)$ satisfying $\lim_{s \to \infty} \gamma(s) = \infty$.

Proposition 2. Let x_* be the unique critical point of a proper C^1 function $V: M \to \mathbb{R}$ on a connected *n*-dimensional C^{∞} manifold *M*. Assume furthermore that *V* is C^{∞} on $M \setminus \{x_*\}$. Then for each class \mathcal{K}_{∞} function γ that is C^{∞} on $(0, \infty)$, there exists a homeomorphism $T: M \to \mathbb{R}^n$ with $T(x_*) = 0$ such that

 $V \circ T^{-1}(y) = \gamma(||y||).$

In particular this holds for $\gamma(||y||) = ||y||^2/2$.

If $n \neq 5$ then T can be chosen to restrict to a C^{∞} -diffeomorphism $M \setminus \{x_*\} \to \mathbb{R}^n \setminus \{0\}$. Furthermore, in this case there exists a class \mathcal{K}_{∞} function γ which is C^{∞} on $(0, \infty)$ and satisfies $\gamma(s)/\gamma'(s) \geq s$ such that T is C^1 with DT(0) = 0.

CRediT authorship contribution statement

Matthew D. Kvalheim: Writing – review & editing, Writing – original draft, Validation, Methodology, Investigation, Funding acquisition, Formal analysis, Conceptualization. Eduardo D. Sontag: Writing – review & editing, Writing – original draft, Methodology, Investigation, Formal analysis, Conceptualization, Funding acquisition.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This material is based upon work supported by the Air Force Office of Scientific Research under award number FA9550-24-1-0299 to Kvalheim and award number FA9550-21-1-0289 to Sontag.

Data availability

No data was used for the research described in the article.

References

- P. Hartman, A lemma in the theory of structural stability of differential equations, Proc. Amer. Math. Soc. 11 (4) (1960) 610–620.
- [2] D.M. Grobman, Homeomorphism of systems of differential equations, Dokl. Akad. Nauk SSSR 128 (5) (1959) 880–881.
- [3] C. Coleman, Local trajectory equivalence of differential systems, Proc. Amer. Math. Soc. 16 (1965) 890–892.
- [4] C. Coleman, Addendum to: Local trajectory equivalence of differential systems, Proc. Amer. Math. Soc. 17 (1966) 770.
- [5] F.W. Wilson Jr., A reformulation of coleman's conjecture concerning the local conjugacy of topologically hyperbolic singular points, in: The Structure of Attractors in Dynamical Systems (Proc. Conf., North Dakota State Univ., Fargo, N.D., 1977), in: Lecture Notes in Math., vol. 668, Springer, Berlin-New York, 1978, pp. 245–252.
- [6] L. Grüne, E.D. Sontag, F.R. Wirth, Asymptotic stability equals exponential stability, and ISS equals finite energy gain—if you twist your eyes, Systems Control Lett. 38 (2) (1999) 127–134.
- [7] G. Perelman, The entropy formula for the Ricci flow and its geometric applications, 2002, pp. 1–39, ArXiv Preprint Math/0211159.
- [8] G. Perelman, Finite extinction time for the solutions to the Ricci flow on certain three-manifolds, 2003, pp. 1–7, ArXiv Preprint Math/0307245.
- [9] G. Perelman, Ricci flow with surgery on three-manifolds, 2003, pp. 1–22, ArXiv Preprint Math/0303109.
- [10] W. Jongeneel, Asymptotic stability equals exponential stability—while you twist your eyes, 2024, pp. 1–27, arXiv preprint arXiv:2411.03277.
- [11] N. Bhatia, G. Szegö, Stability Theory of Dynamical Systems, Number 161 in Grundlehren der Mathematischen Wissenschaften, Springer-Verlag, 1970.
- [12] V.I. Arnold, Ordinary Differential Equations, The MIT Press, 1973, Translated and Edited by Richard A. Silverman.
- [13] Y. Lan, I. Mezić, Linearization in the large of nonlinear systems and Koopman operator spectrum, Phys. D 242 (2013) 42–53.
- [14] J. Eldering, M. Kvalheim, S. Revzen, Global linearization and fiber bundle structure of invariant manifolds, Nonlinearity 31 (9) (2018) 4202–4245.
- [15] M.D. Kvalheim, S. Revzen, Existence and uniqueness of global Koopman eigenfunctions for stable fixed points and periodic orbits, Phys. D 425 (2021) Paper No. 132959, 20.
- [16] M.O. Williams, I.G. Kevrekidis, C.W. Rowley, A data-driven approximation of the Koopman operator: Extending dynamic mode decomposition, J. Nonlinear Sci. 25 (2015) 1307–1346.
- [17] M. Budišić, R. Mohr, I. Mezić, Applied Koopmanism, Chaos 22 (4) (2012) 047510, 33.
- [18] I. Mezić, Spectrum of the koopman operator, spectral expansions in functional spaces, and state-space geometry, J. Nonlinear Sci. 30 (5) (2020) 2091–2145.
- [19] S.L. Brunton, M. Budišić, E. Kaiser, J.N. Kutz, Modern Koopman theory for dynamical systems, SIAM Rev. 64 (2) (2022) 229–340, 4416982.
- [20] G. Haller, B. Kaszás, Data-driven linearization of dynamical systems, Nonlinear Dynam. 112 (21) (2024) 18639–18663.
- [21] M.-A. Belabbas, X. Chen, A sufficient condition for the super-linearization of polynomial systems, Systems Control Lett. 179 (2023) Paper No. 105588, 6.
- [22] Z. Liu, N. Ozay, E.D. Sontag, On the Non-Existence of Immersions for Systems with Multiple Omega-Limit Sets, vol. 56, IFAC World Congress, 2023, pp. 60–64.
- [23] M.D. Kvalheim, P. Arathoon, Linearizability of flows by embeddings, 2024, pp. 1–20, ArXiv Preprint arXiv:2305.18288v6.
- [24] P. Arathoon, M.D. Kvalheim, Koopman embedding and super-linearization counterexamples with isolated equilibria, 2023, pp. 1–7, arXiv preprint arXiv:2306. 15126.
- [25] Z. Liu, N. Ozay, E.D. Sontag, Properties of immersions for systems with multiple limit sets with implications to learning Koopman embeddings, Automatica 176 (2025) 112226.
- [26] J. Ko, M.-A. Belabbas, Minimum number of observables and system invariants in super-linearization, IEEE Control. Syst. Lett. 8 (2024) 2217–2222.
- [27] T. Bárta, R. Chill, E. Fašangová, Every ordinary differential equation with a strict Lyapunov function is a gradient system, Monatsh. Math. 166 (1) (2012) 57–72.
- [28] E.D. Sontag, Contractive systems with inputs, in: J. Willems, S. Hara, Y. Ohta, H. Fujioka (Eds.), Perspectives in Mathematical System Theory, Control, and Signal Processing, Springer-verlag, 2010, pp. 217–228.

- [29] F.W. Wilson Jr., Smoothing derivatives of functions and applications, Trans. Amer. Math. Soc. 139 (1969) 413–428.
- [30] A. Fathi, P. Pageault, Smoothing Lyapunov functions, Trans. Amer. Math. Soc. 371 (3) (2019) 1677–1700.
- [31] F.W. Wilson Jr., The structure of the level surfaces of a Lyapunov function, J. Differential Equations 3 (1967) 323–329.
- [32] J.W. Milnor, Topology from the Differentiable Viewpoint, in: Princeton Landmarks in Mathematics, Princeton University Press, Princeton, NJ, 1997, p. xii+64, Based on notes by David W. Weaver, Revised reprint of the 1965 original.
- [33] M.W. Hirsch, Differential topology, in: Graduate Texts in Mathematics, vol. 33, Springer-Verlag, New York, 1994, p. x+222, Corrected reprint of the 1976 original.
- [34] J. Morgan, G. Tian, Ricci flow and the Poincaré conjecture, in: Clay Mathematics Monographs, vol. 3, American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2007, p. xlii+521.
- [35] S. Smale, On the structure of manifolds, Amer. J. Math. 84 (1962) 387-399.

- [36] M.H. Freedman, The topology of four-dimensional manifolds, J. Differential Geom. 17 (3) (1982) 357–453.
- [37] M.W. Hirsch, On homotopy spheres of low dimension, in: Differential and Combinatorial Topology (a Symposium in Honor of Marston Morse), Princeton Univ. Press, Princeton, NJ, 1965, pp. 199–204.
- [38] E.D. Sontag, Mathematical control theory. Deterministic finite-dimensional systems, second ed., in: Texts in Applied Mathematics, vol. 6, Springer-Verlag, New York, 1998.
- [39] J.J. Bramburger, Data-Driven Methods for Dynamic Systems, Society for Industrial and Applied Mathematics, Philadelphia, PA, 2024.
- [40] M.D. Kvalheim, E.D. Sontag, Why should autoencoders work? Trans. Mach. Learn. Res. (2024).
- [41] M. Morse, The Calculus of Variations in the Large, in: American Mathematical Society Colloquium Publications, vol. 18, American Mathematical Society, Providence, RI, 1996, p. xii+368, Reprint of the 1932 original. MR 1451874.
- [42] J. Milnor, Morse Theory, in: Annals of Mathematics Studies, No. 51, Princeton University Press, Princeton, NJ, 1963, p. vi+153, Based on lecture notes by M. Spivak and R. Wells.